

UNIT III

Logic Concepts: Introduction, propositional calculus, propositional logic, natural deduction system, axiomatic system, semantic tableau system in propositional logic, resolution refutation in propositional logic, predicate logic

1.1. Propositional Logic Concepts:

- Logic is a study of principles used to
 - distinguish correct from incorrect reasoning.
- Formally it deals with
 - the notion of truth in an abstract sense and is concerned with the principles of valid inferencing.
- A proposition in logic is a declarative statements which are either true or false (but not both) in a given context. For example,
 - “Jack is a male”,
 - "Jack loves Mary" etc.
- Given some propositions to be true in a given context,
 - logic helps in inferencing new proposition, which is also true in the same context.
- Suppose we are given a set of propositions such as
 - “It is hot today" and
 - “If it is hot it will rain", then
 - we can infer that

“It will rain today”.

1.2. Well-formed formula

- Propositional Calculus (PC) is a language of propositions basically refers
 - to set of rules used to combine the propositions to form compound propositions using logical operators often called connectives such as \wedge , \vee , \sim , \rightarrow , \leftrightarrow
 - Well-formed formula is defined as:
-

- An atom is a well-formed formula.
- If α is a well-formed formula, then $\sim\alpha$ is a well-formed formula.
- If α and β are well formed formulae, then $(\alpha \wedge \beta)$, $(\alpha \vee \beta)$, $(\alpha \rightarrow \beta)$, $(\alpha \leftrightarrow \beta)$ are also well-formed formulae.
- A propositional expression is a well-formed formula if and only if it can be obtained by using above conditions.

1.3. Truth Table

- Truth table gives us operational definitions of important logical operators.
 - By using truth table, the truth values of well-formed formulae are calculated.
- Truth table elaborates all possible truth values of a formula.

The meanings of the logical operators are given by the following truth table.

P	Q	$\sim P$	$P \wedge Q$	$P \vee Q$	$P \rightarrow Q$	$P \leftrightarrow Q$
T	T	F	T	T	T	T
T	F	F	F	T	F	F
F	T	T	F	T	T	F
F	F	T	F	F	T	T

1.4. Equivalence Laws:

Commutation

1. $P \wedge Q \cong Q \wedge P$
2. $P \vee Q \cong Q \vee P$

Association

1. $P \wedge (Q \wedge R) \cong (P \wedge Q) \wedge R$
2. $P \vee (Q \vee R) \cong (P \vee Q) \vee R$

Double Negation

$$\sim(\sim P) \cong P$$

Distributive Laws

$$1. \quad P \wedge (Q \vee R) \cong (P \wedge Q) \vee (P \wedge R)$$

$$2. \quad P \vee (Q \wedge R) \cong (P \vee Q) \wedge (P \vee R)$$

De Morgan's Laws

$$1. \quad \sim (P \wedge Q) \cong \sim P \vee \sim Q$$

$$2. \quad \sim (P \vee Q) \cong \sim P \wedge \sim Q$$

Law of Excluded Middle

$$P \vee \sim P \cong T \text{ (true)}$$

Law of Contradiction

$$P \wedge \sim P \cong F \text{ (false)}$$

2. Propositional Logic – PL

- PL deals with
 - the validity, satisfiability and unsatisfiability of a formula
 - derivation of a new formula using equivalence laws.
 - Each row of a truth table for a given formula is called its **interpretation** under which a formula can be true or false.
 - A formula α is called **tautology** if and only
 - if α is true for all interpretations.
 - A formula α is also called **valid** if and only if
 - it is a **tautology**.
 - Let α be a formula and if there exist at least one interpretation for which α is true,
 - then α is said to be **consistent** (satisfiable) i.e., if \exists a model for α , then α is said to be consistent .
 - A formula α is said to be inconsistent (unsatisfiable), if and only if
 - α is always false under all interpretations.
-

- We can translate
 - simple declarative and

conditional (if .. then) natural language sentences into its corresponding propositional formulae.

Example

- Show that " It is humid today and if it is humid then it will rain so it will rain today" is a valid argument.
- **Solution:** Let us symbolize English sentences by propositional atoms as follows:

A : It is humid

B : It will rain

- Formula corresponding to a text:

$$\alpha : ((A \rightarrow B) \wedge A) \rightarrow B$$

- Using truth table approach, one can see that α is true under all four interpretations and hence is valid argument.

Truth Table for $((A \rightarrow B) \wedge A) \rightarrow B$				
A	B	$A \rightarrow B = X$	$X \wedge A = Y$	$Y \rightarrow B$
T	T	T	T	T
T	F	F	F	T
F	T	T	F	T
F	F	T	F	T

- Truth table method for problem solving is
 - simple and straightforward and
 - very good at presenting a survey of all the truth possibilities in a given situation.
 - It is an easy method to evaluate
-

- a consistency, inconsistency or validity of a formula, but the size of truth table grows exponentially.
- Truth table method is good for small values of n .
- For example, if a formula contains n atoms, then the truth table will contain 2^n entries.
 - A formula $\alpha : (P \wedge Q \wedge R) \rightarrow (Q \vee S)$ is **valid** can be proved using truth table.
 - A table of 16 rows is constructed and the truth values of α are computed.
 - Since the truth value of α is true under all 16 interpretations, it is valid.
- It is noticed that if $P \wedge Q \wedge R$ is false, then α is true because of the definition of \rightarrow .
- Since $P \wedge Q \wedge R$ is false for 14 entries out of 16, we are left only with two entries to be tested for which α is true.
 - So in order to prove the validity of a formula, all the entries in the truth table may not be relevant.
- Other methods which are concerned with proofs and deductions of logical formula are as follows:
 - Natural Deductive System
 - Axiomatic System
 - Semantic Tableaux Method
 - Resolution Refutation Method

3. Natural deduction method – ND

- ND is based on the set of few deductive inference rules.
- The name natural deductive system is given because it mimics the pattern of natural reasoning.
- It has about 10 deductive inference rules.

Conventions:

- E for Elimination.
 - $P, P_k, (1 \leq k \leq n)$ are atoms.
-

– α_k , ($1 \leq k \leq n$) and β are formulae.

Natural Deduction Rules:

Rule 1: I- \wedge (Introducing \wedge)

I- \wedge : If P_1, P_2, \dots, P_n then $P_1 \wedge P_2 \wedge \dots \wedge P_n$

Interpretation: If we have hypothesized or proved P_1, P_2, \dots and P_n , then their conjunction $P_1 \wedge P_2 \wedge \dots \wedge P_n$ is also proved or derived.

Rule 2: E- \wedge (Eliminating \wedge)

E- \wedge : If $P_1 \wedge P_2 \wedge \dots \wedge P_n$ then P_i ($1 \leq i \leq n$)

Interpretation: If we have proved $P_1 \wedge P_2 \wedge \dots \wedge P_n$, then any P_i is also proved or derived. This rule shows that \wedge can be eliminated to yield one of its conjuncts.

Rule 3: I- \vee (Introducing \vee)

I- \vee : If P_i ($1 \leq i \leq n$) then $P_1 \vee P_2 \vee \dots \vee P_n$

Interpretation: If any P_i ($1 \leq i \leq n$) is proved, then $P_1 \vee \dots \vee P_n$ is also proved.

Rule 4: E- \vee (Eliminating \vee)

E- \vee : If $P_1 \vee \dots \vee P_n, P_1 \rightarrow P, \dots, P_n \rightarrow P$ then P

Interpretation: If $P_1 \vee \dots \vee P_n, P_1 \rightarrow P, \dots$, and $P_n \rightarrow P$ are proved, then P is proved.

Rule 5: I- \rightarrow (Introducing \rightarrow)

I- \rightarrow : If from $\alpha_1, \dots, \alpha_n$ infer β is proved then $\alpha_1 \wedge \dots \wedge \alpha_n \rightarrow \beta$ is proved

Interpretation: If given $\alpha_1, \alpha_2, \dots$ and α_n to be proved and from these we deduce β then $\alpha_1 \wedge \alpha_2 \wedge \dots \wedge \alpha_n \rightarrow \beta$ is also proved.

Rule 6: E- \rightarrow (Eliminating \rightarrow) - Modus Ponens

E- \rightarrow : If $P_1 \rightarrow P, P_1$ then P

Rule 7: I- \leftrightarrow (Introducing \leftrightarrow)

I- \leftrightarrow : If $P_1 \rightarrow P_2, P_2 \rightarrow P_1$ then $P_1 \leftrightarrow P_2$

Rule 8: E- \leftrightarrow (Elimination \leftrightarrow)

$E-\leftrightarrow$: If $P_1 \leftrightarrow P_2$ then $P_1 \rightarrow P_2, P_2 \rightarrow P_1$

Rule 9: I- \sim (Introducing \sim)

I- \sim : If **from P infer $P_1 \wedge \sim P_1$** is proved then **$\sim P$** is proved

Rule 10: E- \sim (Eliminating \sim)

E- \sim : If **from $\sim P$ infer $P_1 \wedge \sim P_1$** is proved then **P** is proved

- If a formula β is derived / proved from a set of premises / hypotheses $\{ \alpha_1, \dots, \alpha_n \}$,
 - then one can write it as **from $\alpha_1, \dots, \alpha_n$ infer β** .
- In natural deductive system,
 - a theorem to be proved should have a form **from $\alpha_1, \dots, \alpha_n$ infer β** .
- Theorem **infer β** means that
 - there are no premises and β is true under all interpretations i.e., β is a tautology or valid.
- If we assume that $\alpha \rightarrow \beta$ is a premise, then we conclude that β is proved if α is given i.e.,
 - if ‘from α infer β ’ is a theorem then $\alpha \rightarrow \beta$ is concluded.
 - The converse of this is also true.

Deduction Theorem: To prove a formula $\alpha_1 \wedge \alpha_2 \wedge \dots \wedge \alpha_n \rightarrow \beta$, it is sufficient to prove a theorem **from $\alpha_1, \alpha_2, \dots, \alpha_n$ infer β** .

Example1: Prove that $P \wedge (Q \vee R)$ follows from $P \wedge Q$

Solution: This problem is restated in natural deductive system as "**from $P \wedge Q$ infer $P \wedge (Q \vee R)$** ". The formal proof is given as follows:

{Theorem} from $P \wedge Q$ infer $P \wedge (Q \vee R)$

{ premise} $P \wedge Q$ (1)

{ E- \wedge , (1)} P (2)

{ E- \wedge , (1)} Q (3)

$$\{ \text{I-V, (3)} \} \quad Q \vee R \quad (4)$$

$$\{ \text{I-}\Lambda, (2, 4) \} \quad P \wedge (Q \vee R) \quad \text{Conclusion}$$

Example2: Prove the following theorem:

$$\text{infer } ((Q \rightarrow P) \wedge (Q \rightarrow R)) \rightarrow (Q \rightarrow (P \wedge R))$$

Solution:

- In order to prove **infer** $((Q \rightarrow P) \wedge (Q \rightarrow R)) \rightarrow (Q \rightarrow (P \wedge R))$, prove a theorem **from** $\{Q \rightarrow P, Q \rightarrow R\}$ **infer** $Q \rightarrow (P \wedge R)$.
- Further, to prove $Q \rightarrow (P \wedge R)$, prove a sub theorem **from** Q **infer** $P \wedge R$

{Theorem} from $Q \rightarrow P, Q \rightarrow R$ **infer** $Q \rightarrow (P \wedge R)$

$$\{ \text{premise 1} \} \quad Q \rightarrow P \quad (1)$$

$$\{ \text{premise 2} \} \quad Q \rightarrow R \quad (2)$$

$$\{ \text{sub theorem} \} \quad \text{from } Q \text{ infer } P \wedge R \quad (3)$$

$$\{ \text{premise } \} \quad Q \quad (3.1)$$

$$\{ \text{E-} \rightarrow, (1, 3.1) \} \quad P \quad (3.2)$$

$$\{ \text{E-} \rightarrow, (2, 3.1) \} \quad R \quad (3.3)$$

$$\{ \text{I-}\Lambda, (3.2, 3.3) \} \quad P \wedge R \quad (3.4)$$

$$\{ \text{I-} \rightarrow, (3) \} \quad Q \rightarrow (P \wedge R) \quad \text{Conclusion}$$

4. Axiomatic System for Propositional Logic:

- It is based on the set of only three axioms and one rule of deduction.
 - It is minimal in structure but as powerful as the truth table and natural deduction approaches.
 - The proofs of the theorems are often difficult and require a guess in selection of appropriate axiom(s) and rules.
 - These methods basically require forward chaining strategy where we start with the given hypotheses and prove the goal.

Axiom1 (A1): $\alpha \rightarrow (\beta \rightarrow \alpha)$

Axiom2 (A2): $(\alpha \rightarrow (\beta \rightarrow \gamma)) \rightarrow ((\alpha \rightarrow \beta) \rightarrow (\alpha \rightarrow \gamma))$

Axiom3 (A3): $(\sim \alpha \rightarrow \sim \beta) \rightarrow (\beta \rightarrow \alpha)$

Modus Ponens (MP) defined as follows:

Hypotheses: $\alpha \rightarrow \beta$ and α **Consequent:** β

Examples: Establish the following:

1. $\{Q\} \vdash (P \rightarrow Q)$ i.e., $P \rightarrow Q$ is a deductive consequence of $\{Q\}$.

{Hypothesis} Q (1)

{Axiom A1} $Q \rightarrow (P \rightarrow Q)$ (2)

{MP, (1,2)} $P \rightarrow Q$ **proved**

2. $\{P \rightarrow Q, Q \rightarrow R\} \vdash (P \rightarrow R)$ i.e., $P \rightarrow R$ is a deductive consequence of $\{P \rightarrow Q, Q \rightarrow R\}$.

{Hypothesis} $P \rightarrow Q$ (1)

{Hypothesis} $Q \rightarrow R$ (2)

{Axiom A1} $(Q \rightarrow R) \rightarrow (P \rightarrow (Q \rightarrow R))$ (3)

{MP, (2, 3)} $P \rightarrow (Q \rightarrow R)$ (4)

{Axiom A2} $(P \rightarrow (Q \rightarrow R)) \rightarrow$
 $((P \rightarrow Q) \rightarrow (P \rightarrow R))$ (5)

{MP, (4, 5)} $(P \rightarrow Q) \rightarrow (P \rightarrow R)$ (6)

{MP, (1, 6)} $P \rightarrow R$ **proved**

4.1. Deduction Theorems in Axiomatic System

Deduction Theorem:

If Σ is a set of hypotheses and α and β are well-formed formulae, then $\{\Sigma \cup \alpha\} \vdash \beta$ implies $\Sigma \vdash (\alpha \rightarrow \beta)$.

Converse of deduction theorem:

Given $\Sigma \vdash (\alpha \rightarrow \beta)$,

we can prove $\{\Sigma \cup \alpha\} \vdash \beta$.

Useful Tips

1. Given α , we can easily prove $\beta \rightarrow \alpha$ for any well-formed formulae α and β .

2. Useful tip

If $\alpha \rightarrow \beta$ is to be proved, then include α in the set of hypotheses Σ and derive β from the set $\{\Sigma \cup \alpha\}$. Then using deduction theorem, we conclude $\alpha \rightarrow \beta$.

Example: Prove $\sim P \rightarrow (P \rightarrow Q)$ using deduction theorem.

Proof: Prove $\{\sim P\} \vdash (P \rightarrow Q)$ and


$\vdash \sim P \rightarrow (P \rightarrow Q)$ follows from deduction theorem.

5. Semantic Tableaux System in PL


- Earlier approaches require
 - construction of proof of a formula from given set of formulae and are called direct methods.
 - In **semantic tableaux**,
 - the set of rules are applied systematically on a formula or set of formulae to establish its consistency or inconsistency.
 - *Semantic tableau*
 - binary tree constructed by using semantic rules with a formula as a root
 - Assume α and β be any two formulae.
-

5.1. Semantic Tableaux Rules


Rule 1: A tableau for a formula $(\alpha \wedge \beta)$ is constructed by adding both α and β to the same path (branch). This can be represented as follows:




Rule 2: A tableau for a formula $\sim(\alpha \wedge \beta)$ is constructed by adding two alternative paths one containing $\sim\alpha$ and other containing $\sim\beta$.




Rule 3: A tableau for a formula $(\alpha \vee \beta)$ is constructed by adding two new paths one containing α and other containing β .




Rule 4: A tableau for a formula $\sim(\alpha \vee \beta)$ is constructed by adding both $\sim\alpha$ and $\sim\beta$ to the same path. This can be expressed as follows:




Rule 5:



Rule 6:




Rule 7:




Rule 8:

$$\alpha \leftrightarrow \beta \equiv (\alpha \wedge \beta) \vee (\sim\alpha \wedge \sim\beta)$$



Rule 9: $\sim(\alpha \leftrightarrow \beta) \equiv (\alpha \wedge \sim\beta) \vee (\sim\alpha \wedge \beta)$



5.2. Consistency and Inconsistency

- If an atom P and $\sim P$ appear on a same path of a semantic tableau,
 - then inconsistency is indicated and such path is said to be **contradictory** or **closed** (finished) path.
 - Even if one path remains **non contradictory** or **unclosed** (open), then the formula α at the root of a tableau is **consistent**.
- **Contradictory tableau** (or **finished tableau**):
 - It defined to be a tableau in which all the paths are contradictory or closed (finished).
- If a tableau for a formula α at the root is a contradictory tableau,
 - then a formula α is said to be inconsistent.

- Show that $\alpha: (Q \wedge \sim R) \wedge (R \rightarrow P)$ is **consistent** and find its model.


{Tableau root} $(Q \wedge \sim R) \wedge (R \rightarrow P)$ (1)

{Apply rule 1 to 1} $(Q \wedge \sim R)$ (2)

$(R \rightarrow P)$ (3)


{Apply rule 1 to 2} Q

{Apply rule 6 to 3} $\sim R$



- $\{Q = T, R = F\}$ and $\{P = T, Q = T, R = F\}$ are models of α .

- Show that $\alpha : (P \wedge Q \rightarrow R) \wedge (\sim P \rightarrow S) \wedge Q \wedge \sim R \wedge \sim S$ is inconsistent using tableaux method.



- α is inconsistent as we get contradictory tableau.

6. Resolution Refutation in PL

- *Resolution refutation*: Another simple method to prove a formula by contradiction.
- Here negation of goal is added to given set of clauses.
 - If there is a refutation in new set using resolution principle then goal is proved
- During resolution we need to identify two clauses,
 - one with positive atom (P) and other with negative atom ($\sim P$) for the application of resolution rule.
- Resolution is based on modus ponens inference rule.

6.1. Disjunctive & Conjunctive Normal Forms

- *Disjunctive Normal Form (DNF)*: A formula in the form $(L_{11} \wedge \dots \wedge L_{1n}) \vee \dots \vee (L_{m1} \wedge \dots \wedge L_{mk})$, where all L_{ij} are literals.
 - Disjunctive Normal Form is disjunction of conjunctions.
- *Conjunctive Normal Form (CNF)*: A formula in the form $(L_{11} \vee \dots \vee L_{1n}) \wedge \dots \wedge (L_{p1} \vee \dots \vee L_{pm})$, where all L_{ij} are literals.

- CNF is conjunction of disjunctions or
- CNF is conjunction of clauses
- *Clause*: It is a formula of the form $(L_1 \vee \dots \vee L_m)$, where each L_k is a positive or negative atom.

6.2. Conversion of a Formula to its CNF

- Each PL formula can be converted into its equivalent CNF.
- Use following equivalence laws:

$$- P \rightarrow Q \cong \sim P \vee Q$$

$$- P \leftrightarrow Q \cong (P \rightarrow Q) \wedge (Q \rightarrow P)$$

- Double Negation

$$- \sim \sim P \cong P$$

- (De Morgan's law)

$$- \sim (P \wedge Q) \cong \sim P \vee \sim Q$$

$$- \sim (P \vee Q) \cong \sim P \wedge \sim Q$$

- (Distributive law)


$$P \vee (Q \wedge R) \cong (P \vee Q) \wedge (P \vee R)$$

6.3 Resolvent of Clauses

- If two clauses C_1 and C_2 contain a complementary pair of literals $\{L, \sim L\}$,
 - then these clauses may be resolved together by deleting L from C_1 and $\sim L$ from C_2 and constructing a new clause by the disjunction of the remaining literals in C_1 and C_2 .
- The new clause thus generated is called **resolvent** of C_1 and C_2 .
 - Here C_1 and C_2 are called parents of resolved clause.
- Inverted binary tree is generated with the last node (root) of the binary tree to be a resolvent.

This is also called resolution tree.

- Find resolvent of the following clauses:
 - $C_1 = P \vee Q \vee R$; $C_2 = \sim Q \vee W$; $C_3 = P \vee \sim W$
- Inverted Resolution Tree



- $\text{Resolvent}(C_1, C_2, C_3) = P \vee R$

6.4 Logical Consequence

- **Theorem1:** If C is a resolvent of two clauses C_1 and C_2 , then C is a *logical consequence* of $\{C_1, C_2\}$.
 - A deduction of an empty clause (or resolvent as contradiction) from a set S of clauses is called a *resolution refutation* of S .
- **Theorem2:** Let S be a set of clauses. A clause C is a *logical consequence* of S iff the set $S' = S \cup \{\sim C\}$ is *unsatisfiable*.
 - In other words, C is a logical consequence of a given set S iff an empty clause is deduced from the set S' .

- Show that $C \vee D$ is a logical consequence of
 - $S = \{A \vee B, \sim A \vee D, C \vee \sim B\}$ using resolution refutation principle.
- First we will add negation of logical consequence
 - i.e., $\sim (C \vee D) \equiv \sim C \wedge \sim D$ to the set S .
 - Get $S' = \{A \vee B, \sim A \vee D, C \vee \sim B, \sim C, \sim D\}$.
- Now show that S' is unsatisfiable by deriving contradiction using resolution principle.

